



**The cyst-theca relation of *Trinovantedinium pallidifulum*,  
with erection of *Protoperidinium lousianensis* sp. nov. and  
their phylogenetic position within the Conica group**

Journal:	<i>Palynology</i>
Manuscript ID	TPAL-2015-0077.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Mertens, Kenneth; Ghent University, Palaeontology; Ifremer, LER BO Gu, Haifeng; Third Institute of Oceanography, SOA Takano, Yoshihito; Institute for East China Sea Research (ECSER) Price, Andrea; McGill University, Geography Pospelova, Vera ; University of Victoria, School of Earth and Ocean Sciences Bogus, Kara; International Ocean Discovery Program, Texas A&M University Versteegh, Gerard; MARUM – Center for Marine Environmental Sciences; Alfred-Wegener-Institut (AWI), Helmholtz Zentrum für Polar und Meeresforschung Marret-Davies, Fabienne ; School of Environmental Sciences, University of Liverpool Turner, R. ; Louisiana State University, Department of Oceanography and Coastal Sciences, Rabalais, Nancy; Louisiana Universities Marine Consortium Matsuoka, Kazumi; Institute for East China Sea Research (ECSER)
Keywords:	Gulf of Mexico, Saanich Inlet, microFTIR, Selenopemphix undulata, Wadden Sea, lake Saroma

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**The cyst-theca relation of *Trinovantedinium pallidifulum*, with erection of *Protoperidinium louisianensis* sp. nov. and their phylogenetic position within the *Conica* group**

Kenneth Neil Mertens<sup>1,\*</sup>, Haifeng Gu<sup>2</sup>, Yoshihito Takano<sup>3</sup>, Andrea M. Price<sup>4</sup>, Vera Pospelova<sup>5</sup>, Kara Bogus<sup>6</sup>, Gerard J. M. Versteegh<sup>7,†</sup>, Fabienne Marret<sup>8</sup>, R. Eugene Turner<sup>9</sup>, Nancy N. Rabalais<sup>10</sup>, Kazumi Matsuoka<sup>3</sup>

<sup>1</sup>Research Unit for Palaeontology, Ghent University, Krijgslaan 281 s8, 9000 Ghent, Belgium

<sup>2</sup>Third Institute of Oceanography, SOA, Xiamen 361005, China, guhaifeng@tio.org.cn

<sup>3</sup>Institute for East China Sea Research (ECSER), 1551-7, Taira-machi, Nagasaki, 851-2213, Japan, takadino@gmail.com, kazu-mtk@nagasaki-u.ac.jp

<sup>4</sup>Department of Geography, McGill University, Burnside Hall, 805 Sherbrooke Street West, Montreal, QC, Canada H3A OB9, andrea.price@mail.mcgill.ca

<sup>5</sup>School of Earth and Ocean Sciences, University of Victoria, OEASB A405, P.O. Box 1700 STN CSC, Victoria, British Columbia, V8W 2Y2, Canada, vpospe@uvic.ca

<sup>6</sup>International Ocean Discovery Program, Texas A&M University, 1000 Discovery Drive, College Station, TX, USA 77845, bogus@iodp.tamu.edu

<sup>7</sup>MARUM – Center for Marine Environmental Sciences, Leobener Strasse, 28359 Bremen, Germany, versteegh@uni-bremen.de

<sup>8</sup>School of Environmental Sciences, University of Liverpool, Liverpool, L69 7ZT, UK f.marret@liv.ac.uk

<sup>9</sup>Department of Oceanography and Coastal Sciences, Louisiana State University, Baton Rouge, Louisiana 70803 USA, euturne@lsu.edu

<sup>10</sup>Louisiana Universities Marine Consortium, Chauvin, Louisiana 70344 USA, nrabalais@lumcon.edu

We establish the cyst-theca relationship of the cyst species *Trinovantedinium pallidifulum* Matsuoka 1987 based on germination experiments of specimens isolated from the Gulf of Mexico. We show that the motile stage is a new species, designated as *Protoperidinium louisianensis*. We also determine its phylogenetic position based on single-cell PCR of a single cell germinated from the Gulf of Mexico cysts. To further refine the phylogeny, we determined the LSU sequence through single-cell PCR of the cyst *Selenopemphix undulata* isolated from Brentwood Bay (Saanich Inlet, BC, Canada). The phylogeny shows that *P. louisianensis* is closest to *P. shanghaiense*, the motile stage of *T. applanatum*, and ~~leave the possibility is consistent with that the monophyly of the genus *Trinovantedinium* is monophyletic~~. *Selenopemphix undulata* belongs to a different clade than *Selenopemphix quanta* (alleged cyst of *P. conicum*), suggesting that the genus *Selenopemphix* is polyphyletic. *Trinovantedinium pallidifulum* is widely distributed with occurrences in the Gulf of Mexico, the North Atlantic, the NE Pacific and SE Asia. In addition, we illustrate the two other extant species, *Trinovantedinium applanatum* and *Trinovantedinium variabile* and two morphotypes of *Trinovantedinium*. Geochemical analyses of the cyst wall of *T. pallidifulum* indicate the presence of amide groups in

\* Now at: Ifremer, LER BO, Station de Biologie Marine, Place de la Croix, BP40537, F-29185 Concarneau Cedex, France, kenneth.mertens@ifremer.fr

† Corresponding author. Email: kenneth.mertens@ifremer.fr

‡ Now at: Alfred-Wegener-Institut (AWI), Helmholtz Zentrum für Polar und Meeresforschung, Am Handelshafen 12, D-27570, Bremerhaven, Germany, gerard.versteegh@awi.de

agreement with other heterotrophic dinoflagellate species, although the cyst wall cyst of *T. pallidifulum* also includes some unique features.

**Keywords:** microFTIR; *Selenopemphix undulata*; Wadden Sea; lake Saroma; Saanich Inlet; Gulf of Mexico

## 1. Introduction

The heterotrophic thecate dinoflagellate genus *Protoperidinium* Bergh is characterized by possessing three cingular plates and an additional transitional plate (Balech 1974). This large genus currently encompasses approximately 280 species (Gómez 2012). The identification of the thecate stage of *Protoperidinium* species is based on the body size, outline, presence and position of apical and/or antapical horns/spines, cingulum displacement and particularly the plate pattern (e.g., Hoppenrath et al. 2009). In regard to the thecal plate arrangement, Balech (1974) used the number of anterior intercalary plates and precingular plates to subdivide *Protoperidinium* into three subgenera: the subgenus *Protoperidinium*, which has seven precingular plates and three anterior intercalary plates; the subgenus *Minusculum*, which has six precingular plates and three anterior intercalary plates; and the subgenus *Archaeoperidinium*, which has seven precingular plates and two anterior intercalary plates. Later, the subgenus *Testeria* was erected by Faust (2006) to accommodate species with seven precingular plates, one anterior intercalary plate and no apical pore complex. However, the results from molecular phylogeny questioned the validity of the subgenus *Minusculum* because it was nested within the subgenus *Protoperidinium* (Yamaguchi et al. 2007; Ribeiro et al. 2010), and showed the polyphyly of the subgenus *Archaeoperidinium*, as originally described by Jörgensen (1912) (Ribeiro et al. 2010). *Archaeoperidinium* was, therefore, emended as a genus and characterized by the flat sulcus, sulcal flagellar fin covering the sulcal area and circular cingulum without displacement (Yamaguchi et al. 2011).

The subgenus *Protoperidinium* can be subdivided into several sections based on the shape of the first apical (1') plate (ortho, meta or para) and the shape of the second anterior intercalary (2a) plate (quadra, penta or hexa) (e.g., Gribble and Anderson 2006). Molecular phylogenies suggested that most of these sections are monophyletic and nested within the *Protoperidinium* sensu stricto clade (Mertens et al. 2013; Gu et al. 2015). There are only two exceptions: the section *Conica* is polyphyletic and positioned within the *Protoperidinium* sensu stricto clade (Yamaguchi et al. 2006; Gu et al. 2015), and the section *Oceanica* is monophyletic but positioned outside of the *Protoperidinium* sensu stricto clade (Sarai et al. 2013).

Several *Protoperidinium* species have been associated ~~with~~ a particular cyst, and the morphology of these cysts can be taxonomically informative, especially the shape of ~~the opening in the cyst wall, termed~~ the archeopyle (Harland 1982). Thecate stages belonging to *Protoperidinium* and *Archaeoperidinium* with a transitional plate ~~have cysts~~ generally ~~havewith~~ a saphopylic or theropylic 2a archeopyle ~~of their respective cyst~~ (Harland 1982; Ribeiro et al. 2010; Mertens et al. 2012a) whereas, *Protoperidinium* species without a transitional plate have ~~cysts with~~ a compound archeopyle involving 2'–4' (Lewis & Dodge 1987; Kawami & Matsuoka 2009; Kawami et al. 2009; Mertens et al. 2013; Liu et al. 2014). Cyst morphology can be diverse and is used to classify cysts into various cyst-~~defined~~ genera such as *Brigantedinium* Reid, *Votadinium* Reid and *Selenopemphix* Benedek (Fensome et al. 1993). However, differences between the cyst-~~based-defined~~ nomenclature and the motile-~~based-defined~~ nomenclature have not yet been fully reconciled.

1  
2  
3  
4  
5  
6  
7  
8  
9 89 | *Trinovantedinium* Reid is a cyst-based-defined genus erected by Reid (1977) to describe  
10 90 peridiniacean cysts with an intercalary archeopyle because, at that time, the existence of such an  
11 91 archeopyle in the genus *Lejeunecysta* Artzner & Dörhöfer 1978 was in doubt. This initial  
12 92 diagnosis included species with or without processes on the cyst. The genus was subsequently  
13 93 emended by Harland (1977), Bujak (1984) and de Verteuil & Norris (1992). The latter restricted  
14 94 the diagnosis of *Trinovantedinium* to only include biphragmal cysts with short, penitabular and  
15 95 intratabular, but never sutural, hollow to solid processes. There are currently 11 species within  
16 96 this genus (Table 1). The type species was described by Reid (1977) and is the large transparent  
17 97 *Trinovantedinium capitatum* from recent-modern Rosslare Point sediment (Ireland). Later, this  
18 98 was considered a junior synonym of *Lejeunia applanata*, described by Bradford (1977) from  
19 99 recent sediments off the east coast of the Musandam Peninsula (Oman), which was renamed later  
20 100 as *Trinovantedinium applanatum* (for details see de Verteuil & Norris 1992, p. 408). This is the  
21 101 first of only three species from this genus that are considered extant; the other eight species are  
22 102 extinct (Table 1). *Trinovantedinium pallidifulum* is the second extant species, first described by  
23 103 Matsuoka (1987) from Holocene surface sediments in Akkeshi Bay (Hokkaido, Japan). The third  
24 104 is *Trinovantedinium variabile*, described by Bujak (1984) from the early Pliocene of the Bering  
25 105 Sea, and was recorded as extant by Radi & de Vernal (2004), Pospelova et al. (2008),  
26 106 Krepakevich & Pospelova (2010), Price & Pospelova (2011) and Bringué et al. (2013).  
27 107 For only one of these extant species is the cyst-theca relationship and molecular  
28 108 phylogenetic position known. The cyst-theca relationship for the type species *Trinovantedinium*  
29 109 *applanatum* was first established by Wall & Dale (1968); however, it was mistakenly related to  
30 110 *Protoperidinium pentagonum* (Gran) Balech. This error was propagated in subsequent studies  
31 111 (Matsuoka 1982; Lewis et al. 1984; Baldwin 1987). It was Inoue, H. in Fukuyo et al. (1990, p.  
32 112 154–155) who first remarked that the cingulum of the *T. applanatum* thecate stage has no  
33 113 displacement, in contrast with *P. pentagonum*. Gu et al. (2015) re-established the cyst-theca  
34 114 relationship for *T. applanatum* from the East China Sea, and they erected the species  
35 115 *Protoperidinium shanghaiense* to describe the motile stage. The hexa-ortho configuration placed  
36 116 this species in the *Conica* group and its LSU rDNA sequence was closest to *P. divaricatum*, *P.*  
37 117 *leonis* and *P. conicum*. This cyst-theca relationship was most recently confirmed by Li et al.  
38 118 (2015).  
39 119 In this study, we establish the cyst-theca relationship for the second extant species of the  
40 120 genus *Trinovantedinium*, *T. pallidifulum*, through incubation of surface sediments from the Gulf  
41 121 of Mexico, Dee Estuary (United Kingdom), and Wadden Sea (Germany). We erect a new  
42 122 species, *Protoperidinium louisianensis*, to describe the motile stage of *Trinovantedinium*  
43 123 *pallidifulum*. We obtained LSU rDNA sequences through single-cell PCR that show the motile  
44 124 stage is closest to *Protoperidinium shanghaiense*. In addition, we document the distribution of  
45 125 *Trinovantedinium pallidifulum* and the geochemical composition of its cyst wall. We also  
46 126 illustrate the two other extant species, *Trinovantedinium applanatum* and *Trinovantedinium*  
47 127 *variabile*. Finally, to further constrain the phylogeny, we also resolve the phylogenetic position  
48 128 of *Selenopemphix undulata* from Brentwood Bay (B.C., Canada).  
49 129  
50 130 **2. Material and methods**  
51 131 **2.1. Germination experiments**  
52 132 We collected cysts of *Trinovantedinium pallidifulum* for incubation studies from surface  
53 133 sediment samples at three locations: (1) northern Gulf of Mexico, (2) Dee Estuary (UK), and (3)  
54 134 the Wadden Sea in northern Germany (Figure 1 and Table 2). All samples were stored in plastic

bags and refrigerated at 4 °C. In situ sea-surface salinities (SSSs) and sea-surface temperatures (SSTs) were measured when collecting the samples (Table 2).

Approximately 0.5–1.0 cm<sup>3</sup> of wet sediment was immersed in filtered seawater after which it was ultrasonicated in a bath (60 s) and rinsed through a 20 µm nylon mesh using filtered seawater. The cyst fraction was separated from this residue using heavy liquid sodium polytungstate (density = 1.3 g cm<sup>-3</sup>) (Bolch 1997). Single cysts were then transferred to Nunclon 0.5 ml microwells subjected to an irradiance of 100 µmol photons m<sup>-2</sup> s<sup>-1</sup> and 24 h light, and filled with L1 medium. The wells were kept at room temperature. Cysts were regularly checked for germination, and observations were performed under an Leitz DM inverted light microscope. Encysted and excysted cysts as well as motile cells were photographed and measured using an Leica DM 5000B light microscope equipped with a Leica DFC 490 camera with 100x oil immersion objectives. For each motile cell, the length, width, depth, distance between the tips of the antapical horns, and width of the cingulum were measured, as where possible. For each cyst, the same parameters were measured; additionally, the length of three randomly chosen spines per cyst were measured. All measurements in the species descriptions cite, in order: the minimum, average (in parentheses) and maximum values (in µm). The standard deviation (SD) is also provided where appropriate. Incubation experiments were done by KNM and took place at GEOTOP (Gulf of Mexico and Dee Estuary samples) and Univ. of Bremen (Wadden Sea samples). All measurements were done by KNM.

We also attempted to germinate *Selenopemphix undulata* from Saanich Inlet (Canada), but none of the isolated cysts germinated.

## 2.2. Study of cysts from surface sediments

To determine the distribution of *Trinovantedinium pallidifulum*, permanent slides of surface samples were examined which included locations from the the northeastern and northwestern Pacific and the northern Gulf of Mexico (Figure 1 and Table 2). Routine palynological techniques were used for processing (Pospelova et al. 2004; Matsuoka et al. 2003; Mertens et al. 2012b). The samples were oven-dried at 40 °C and then treated with room-temperature 10% hydrochloric acid (HCl) to remove calcium carbonate. The material was rinsed twice with distilled water, sieved at 120 µm to eliminate the coarse fraction, and retained on a 15 µm nylon mesh. To dissolve siliceous particles, samples were treated with 48–50% room-temperature hydrofluoric acid (HF) for at least two days, and then treated for 10 min with room-temperature HCl (10%) to remove fluorosilicates. The residue was rinsed twice with distilled water, ultrasonicated for 30 s and finally collected on a 15 µm mesh. Residue aliquots were mounted in glycerine jelly. All measurements and light photomicrographs were as described in Section 2.1. In addition, in order to illustrate *Trinovantedinium applanatum* and *Trinovantedinium variabile*, we re-examined permanent slides from palynologically prepared samples from several localities.

## 2.3. Single-cell PCR amplification and sequencing of the motile stage of *Trinovantedinium pallidifulum*

Surface sediment samples containing *Trinovantedinium pallidifulum* were used from the Gulf of Mexico (Figure 1 and Table 2). Cysts were isolated from the sediment using the heavy liquid separation described in Section 2.1. Motile cells identified through light microscopy were rinsed several times in sterilized distilled water, broken by compressing the cell between the slide and cover slip, and then transferred into a PCR tube. The single cell was used as the template to amplify about 1200 bp of the nuclear-encoded LSU rDNA, using the primers D1R (Scholin et al.



1  
2  
3  
4  
5  
6  
7  
8  
9 181 1994) and 28-1483R (Daugbjerg et al. 2000). A 50 µl PCR cocktail containing 0.2 µM primers,  
10 182 PCR buffer, 50 µM dNTP mixture, 1U of Ex Taq DNA polymerase (Takara, Dalian, China) was  
11 183 subjected to 35 cycles using a Mastercycler PCR (Eppendorf, Hamburg, Germany). The PCR  
12 184 protocol was: initial denaturation for 3.5 min at 94 °C, followed by 35 cycles of 50 s denaturation  
13 185 at 94 °C, 50 s annealing at 45 °C, and 80 s extension at 72 °C, plus a final extension of 10 min at  
14 186 72 °C. The amplified products were run on a 1% agarose gel. Positive bands were excised and  
15 187 purified using a DNA extraction kit (Sangon, Shanghai, China) and sequenced in both directions  
16 188 using the ABI Big-Dye dye-terminator technique (Applied Biosystems, Foster City, California,  
17 189 USA), according to the manufacturer recommendations. This DNA work was performed by HG  
18 190 in Third Oceanographic Centre, Xiamen, China.  
19 191

19 192 **2.4. Single-cell PCR amplification and sequencing of cysts of *Selenopemphix undulata***  
20 193 We obtained surface sediment samples containing *Selenopemphix undulata* from Brentwood Bay  
21 194 (B.C., Canada) (48,58°N,-123.47°E, 6 m water depth) using a petite ponar grab on October 1,  
22 195 2011. Cysts were isolated from the sediment using heavy liquid separation as described in  
23 196 Section 2.1. The cysts were then rinsed several times in sterilized distilled water, broken by  
24 197 compressing the cell between the slide and cover slip, and then transferred into a PCR tube. The  
25 198 single cell was used as the template to amplify about 1200 bp of the nuclear-encoded LSU  
26 199 rDNA, using the primers D1R (Scholin et al. 1994) and 28-1483R (Daugbjerg et al. 2000). A 50  
27 200 µl PCR cocktail containing 0.2 µM primers, PCR buffer, 50 µM dNTP mixture, 1U of Ex Taq  
28 201 DNA polymerase (Takara, Dalian, China) was subjected to 35 cycles using a Mastercycler PCR  
29 202 (Eppendorf, Hamburg, Germany). The PCR protocol was an initial denaturation for 3.5 min at 94  
30 203 °C, followed by 35 cycles of 50 s denaturation at 94 °C, 50 s annealing at 45 °C, and 80 s  
31 204 extension at 72 °C, plus a final extension of 10 min at 72 °C. The amplified products were run on  
32 205 a 1% agarose gel. Positive bands were excised and purified using a DNA extraction kit (Sangon,  
33 206 Shanghai, China) and sequenced in both directions using the ABI Big-Dye dye-terminator  
34 207 technique (Applied Biosystems, Foster City, California, USA), according to the manufacturer's  
35 208 recommendations. This DNA work was performed by YT at the University of Nagasaki, Japan.  
36 209

36 210 **2.5. Sequence alignments and phylogenetic analyses**  
37 211 Newly obtained sequences were first aligned with those of related species available in GenBank  
38 212 using 'BioEdit' v7.0.0 (Hall 1999), and subsequently using Mafft (Katoh et al. 2005)  
39 213 (<http://mafft.cbrc.jp/alignment/server/>). *Akashiwo sanguinea* (Hirasaka) G. Hansen & Moestrup  
40 214 was selected as the outgroup. A Bayesian reconstruction of the data matrix was performed with  
41 215 MrBayes 3.0b4 (Ronquist & Huelsenbeck 2003) using a general time reversible model (GTR  
42 216 +I+G) chosen by JmodelTest (Posada 2008). Four Markov chain Monte Carlo (MCMC) chains  
43 217 ran for two million generations, sampling every 1,000 generations with a burnin of 10%. A  
44 218 majority rule consensus tree was created in order to examine the posterior probabilities of each  
45 219 clade. Maximum likelihood-based analyses were conducted with RaxML v7.2.6 (Stamatakis  
46 220 2006) on the T-REX web server (Boc et al. 2012) using the above model. Bootstrap values were  
47 221 determined with 1,000 replicates.  
48 222

48 223 **2.6. Geochemical analysis of cyst wall chemistry**  
49 224 After germination, an empty cyst originally derived from surface sediment of the Wadden Sea  
50 225 (Germany) (Figure 1) was removed from a microwell using a micropipette into a droplet of water  
51 226 contained on a glass slide with a concave depression. The water was allowed to evaporate and  
52  
53  
54  
55  
56  
57  
58  
59  
60

then a droplet of ethanol was added. The cyst was allowed to soak in the ethanol for 30 min after which a droplet of MilliQ water was added. The cyst was then isolated and the procedure repeated twice. These steps were to ensure that any polar or apolar compounds adhered to the cyst wall were removed. The cyst was then visually examined under the light microscope and placed on an Au-coated mirror and analysed with micro-Fourier transform infrared spectroscopy using a Bruker FT-IR microscope (Hyperion 3000) with a 15x objective. The spectrum was acquired in reflective mode with 50 scans over 4000–600 cm<sup>-1</sup> and is shown after background subtraction. Peak assignments were based on Colthup (1990) and relevant published literature (e.g., Bogus et al. 2012; 2014; Cárdenas et al. 2004; Versteegh et al. 2012). [This geochemical work was done by KB, KNM and GJMV at the Univ. of Bremen, Germany.](#)

### 3. Results

#### 3.1. Results of germination experiments

Undescribed motile cells, here assigned to *Protoperidinium louisianensis* n. sp., emerged from *Trinovantedinium pallidifulum* cysts isolated from surface sediments of the Gulf of Mexico (five specimens identified) (Figure 1 and Table 1). [These motile cells germinated from the cysts](#) After one or two days of incubation, ~~motile cells germinated from the cysts~~. These cells died a few days after germination and never divided. Two out of four specimens of *Trinovantedinium pallidifulum* from the Wadden Sea (Germany) germinated, but the cells did not fully develop thecal plates. A single specimen from [the](#) Dee Estuary (UK) germinated, but the motile stage could not be fully observed.

#### 3.2. Systematic palaeontology

Division DINOFLAGELLATA (Bütschli 1885) ~~emend.~~ Fensome et al. 1993, emend. Adl et al. 2005

Class DINOPHYCEAE Pascher 1914

Subclass PERIDINIPHYCIDA Fensome et al. 1993

Order PERIDINIALES Haeckel 1894

Family PROTOPERIDINIACEAE Balech 1988 nom. cons.

Subfamily PROTOPERIDINIOIDEAE (Autonym)

Genus *Protoperidinium* Bergh 1881

*Protoperidinium louisianensis* Mertens, Gu, Price et Matsuoka n. sp.

Plate 1, figures 1–15, Plate 2, figures 1–15

**Holotype.** Plate 1, figures 1–9.

**Type locality.** Northern Gulf of Mexico, station A7 (28.94°N, 89.75°W), offshore Louisiana.

**Diagnosis.** A species of intermediate size of the genus *Protoperidinium* with the tabulation formula Po, X, 4', 3a, 7'', 3c+t, ?s, 5''', 2'''. The motile cell is pentagonal in outline and dorsoventrally flattened, with a short apical horn and two antapical horns, each bearing a short spine. The epitheca is longer than the hypotheca, and bears convex sides. Plate 1' is ortho-type, 1a and 3a are penta-type, and 2a is hexa-type and stenodeltaform linteloid. Plates are thin with polygonal reticulations. The cyst is pentagonal and light brown in color, with a thickened apical horn and two thickened antapical horns. The cyst surface is smooth, bearing numerous peritabular, short, solid, erect, and non-branching processes with acuminate tips. Sometimes the dorsal side of the hypotheca is striated. The archeopyle is stenodeltaform linteloid, angular and saphopylic.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Derivation of name.** The specific epithet refers to the type locality, which liesis offshore the state of Louisiana (USA).

**Gene sequence.** The LSU rDNA gene sequence of the cyst—GenBank Accession No. KU519754XXXXXX (LSU).

**Description.** *Description of motile cell of* *Protoperidinium louisianensis* (Plate 1, figures 1–9, Plate 2, figures 1–9). The excysted motile cells (five observed and not preserved) were pentagonal in outline, dorsoventrally flattened and carried an apical horn and two antapical horns of equal length, each bearing a short spine (Plate 1, figure 1–2, Plate 2, figure 1). The epitheca had convex sides and was longer than the hypotheca. The cell contents were greenish, except for a red body. The thin thecal plates carried polygonal reticulations (Plate 1, figure 2,7).

The plate arrangement on the epitheca was bilaterally symmetrical. The oval apical pore plate (Po) was surrounded by a low apical collar formed by the raised edges of the apical plates (Plate 2, figure 4). The canal plate (X) was elongate and trapezoidal (Plate 2, figure 4). The first apical plate (1') was wide and rhombic (ortho-type) and the sides of plate 1' contacting plates 2' and 4' are longer than those contacting plates 1'' and 7'' (Plate 1, figure 2). Plates 2' and 4' were elongated and subpentagonal, whereas 3' was short and subpentagonal (Plate 1, figure 4). The first and third anterior intercalary plates (1a) were pentagonal and equal in size (Plate 1, figures 3,6). The second anterior intercalary plate (2a) was hexagonal, stenodeltaform linteloid and more elongated and had two small sides touching plates 3'' and 5'' (Plate 1, figure 4). The precingular series consisted of seven plates. Plate 1'', 4'' and 7'' were quadrangular (Plate 1, figures 2,4), and 2'', 3'', 5'' and 6'' pentagonal (Plate 1, figures 3,5,6). The cingulum was slightly left-handed (descending), lined with narrow lists and comprising three cingular plates plus a transitional plate. The transitional plate (t) was small. Plate 1c reached the end of plate 1'' and 2''' (Plate 1, figure 3). Plate 2c was the longest of the series and reached a short way beyond the 6''/7'' boundary and the 4'''/5''' boundary (Plate 1, figure 7). Plate 3c was similar in size to Plate 1c.

We were unable to dissect and observe all the sulcal plates.

The plate arrangement of the hypotheca was also symmetrical, featuring five postcingular plates. Plate 5''' was longer than plate 1'''. Plates 1''', 3''', and 5''' were pentagonal, and 2''' and 4''' were quadrangular (Plate 1, figures 7–9). The antapical series comprised two plates, 1'''' and 2'''', which formed the antapical horns (Plate 1, figure 7).

The plate formula is thus Po, X, 4', 3a, 7'', 3c+t, ?s, 5''', 2''''', and the complete tabulation (except for the sulcal plates) is illustrated in Figure 2.

*Description of cyst of* *Protoperidinium louisianensis* (Plate 1, figures 10–15, Plate 2, figures 10–15). Cysts were similar in shape to the motile stage, but light brown in color, bearing numerous small solid spines. The cysts were peridinioid (pentagonal) with a thickened apical horn and two thickened antapical horns of equal length. Living cysts contained abundant greenish granules. The epicyst had convex sides and was always longer than the hypocyst. The central body wall was thin (>0.3 μm) and biphragmal with closely appressed layers that separate along the apical horn and antapical horns, with a smooth surface (Plate 1, figure 11). Sometimes striations were present on the dorsal side of the hypocyst (Plate 2, figure 15). The processes were short, solid, erect, and non-branching with acuminate tips (Plate 1, figure 11). The process distribution was largely peritabular-penitabular (Plate 1, figure 13, Plate 2, figure 11). The process length was fairly constant for individual specimens, except around the apical horn and antapical horns where they became longer (Plate 2, figure 10). The paracingulum was excavated with lists ornamented with rows of equidistant processes, and slight left-handed displacement (Plate 2, figure 11). The parasulcus was free of processes and indented (Plate 1, figure 14), two flagellar scars were

Formatted: Font: Times New Roman, 12 pt



always visible (Plate 2, figure 12). The archeopyle was angular and saphopylic, involving release of plate 2a, steno-deltaform linteloid. The description is based on cysts used in the incubation experiments and those recovered from sediment and prepared using palynological methods. Cysts without cell content in the palynologically treated samples did not retain their shape well - they did not seem to be very robust as they were always quite folded and flattened.

**Dimensions.** Incubated motile cells: length, 52.3 (58.8) 65.1  $\mu\text{m}$  (SD = 6.0, n=4); width, 44.0 (53.6) 58.8  $\mu\text{m}$  (SD = 7.4, n=4); depth, 34.8 (38.4) 41.9  $\mu\text{m}$  (SD = 5.0, n=2); distance between the tips of the antapical horns, 12.8 (16.3) 20.3  $\mu\text{m}$  (SD = 3.7, n=4); width of cingulum, 3.9 (4.1) 4.4  $\mu\text{m}$  (SD = 0.2, n=4).

Cysts germinated to give identifiable thecae: length, 43.0 (58.5) 68.4  $\mu\text{m}$  (SD = 8.2, n=10); width, 46.3 (56.5) 63.9  $\mu\text{m}$  (SD = 5.0, n=10); depth, 30.4 (40.6) 48.4  $\mu\text{m}$  (SD = 7.2, n=7); distance between the tips of the antapical horns, 16.0 (23.9) 27.7  $\mu\text{m}$  (SD = 3.6, n=10); width of cingulum, 4.5 (6.0) 7.2  $\mu\text{m}$  (SD = 0.8, n=10); average length of three spines per cyst, 1.1 (1.8) 2.7  $\mu\text{m}$  (SD = 0.4, n=30).

Cysts palynologically prepared from surface sediments of different locations: length, 49.9 (56.6) 62.8  $\mu\text{m}$  (SD = 5.7, n=5); width, 49.8 (56.3) 63.3  $\mu\text{m}$  (SD = 5.0, n=5); depth, none measured; distance between the tips of the antapical horns, 20.6 (21.3) 22.5  $\mu\text{m}$  (SD = 0.8, n=5); width of cingulum, 4.8 (5.6) 6.5  $\mu\text{m}$  (SD = 0.7, n=4); average length of three spines per cyst, 1.4 (1.9) 2.8  $\mu\text{m}$  (SD = 0.4, n=15).

**Comments.** The geological preservability of the cysts was demonstrated by their ability to withstand palynological treatment and presence in sediments at least as old as Holocene (Matsuoka et al. 1999) to mid-Miocene (our interpretation of the "undefined protoperidiniacean species" depicted in de Verteuil & Norris (1992), plate 2, figs. 9–12). Specimens from the German Wadden Sea show an identical morphology (Plate 3, figures 1–12) and the cysts correspond to the cyst-based-defined species *Trinovantedinium pallidifulum* Matsuoka, here reillustrated by its holotype (Plate 3, figures 13–16).

Genus *Trinovantedinium* Reid 1977, emend. de Verteuil & Norris 1992

*Trinovantedinium applanatum* (Bradford 1977) Bujak & Davies 1983

Plate 4, figures 1–13, Plate 5, figures 1–10

**Synonyms.** *Trinovantedinium capitatum* Reid 1977, Plate 1, figures 6–8.

**Comments.** The illustrated specimens conform to the original description of Bradford (1977). We also illustrate two extreme morphotypes of this species. Type A (Plate 5, figures 1–7) was found in warmer water regions (Omura Bay, Japan; Bay of Bengal; Red Sea) and shows a more elongate, transparent body shape with straight sides and more elongated horns, separated by a deep depression. Type B (Plate 5, figures 8–10) was found in cold water (offshore Greenland) and shows a more rounded, transparent body shape with reduced horns and a very shallow depression between the antapical horns. *Trinovantedinium applanatum* differs from *Trinovantedinium pallidifulum* in its transparency, often larger and more elongate body and longer processes.

*Trinovantedinium variabile* (Bujak 1984) de Verteuil & Norris 1992

Plate 6, figures 1–11

**Comments.** The illustrated specimens conform to the original description of Bujak (1984). The holotype is redescribed and reillustrated by Head (1994, p. 226, pl. 11, figs. 4, 5, 7, 8).

Formatted: French (France)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Genus *Selenopemphix* Benedek 1972, emend. Head 1993  
*Selenopemphix undulata* Verleye, Pospelova, Mertens et Louwye 2011, [Plate 1, figure 8](#)  
Plate 6, figures 12–14

**Comments.** One cyst corresponding to *Selenopemphix undulata* as described by Verleye et al. (2011, [particularly the specimen displayed in Plate 1, figure 8](#)) was isolated for single-cyst PCR (Plate 6, figures 12–14). The cyst was collected from surface sediment of Brentwood Bay (B.C., Canada). The cyst was large and polar compressed, with a cingulum with [undulating](#)-margins [showing very small undulations](#). The epicyst was conical and striated, the hypocyst was also striated and bore two fused antapical horns that were wider than the apical horn. The archeopyle could not be observed on encysted specimens. The single cyst had a length of 44.5 µm (apex to antapex), width of 74.6 µm, and depth of 70.8 µm.  
**Gene sequence.** The LSU rDNA gene sequence of the cyst—[GenBank Accession No. LC114019 \(LSU\)](#).

**3.3. Phylogenetic position of *Protoperidinium louisianensis* and *Selenopemphix undulata* as inferred from LSU rDNA sequences**

We obtained 1,143 base pairs from one germinated cell of *Protoperidinium louisianensis* isolated from the Gulf of Mexico (Accession number: [KU519754XXXXXX](#)), and this sequence was used for the phylogenetic analyses (Figure 3). *Protoperidinium louisianensis* was closest to *Protoperidinium shanghaiense*, and formed a clade with several other species belonging to the *Conica* section (*P. divaricatum*, *P. conicum*, *P. leonis*). *Selenopemphix undulata* ([Accession number: LC114019](#)), was closest to *Protoperidinium biconicum*, which also belongs to the section *Conica* and formed a separate clade with *P. punctulatum* and *P. humile*, both species belonging to the section *Tabulata*. The other members of the Protoperidiniaceae formed several other clades, which were also used by Gu et al. (2015).

**3.4. ~~Modern Recent~~ distribution of *Trinovantedinium pallidifulum* and inferred ecology**

*Trinovantedinium pallidifulum* was initially described by Matsuoka (1987) from Akkeshi Bay (Hokkaido, Japan) and subsequently predominantly in southeast Asia, [such as including](#) surface sediments off South Korea (Cho et al. 2003; Shin et al. 2011), Japan (Kojima et al., 1994; Matsuoka et al. 2003), China (Wang et al. 2004), and Malaysia (Furio et al. 2006) (Figure 1)).

In this study we show that this species is much more widely distributed since it is recorded in surface sediments from the northern Gulf of Mexico, Casino Coast (Brazil), the North Atlantic (La Vilaine Bay, Wadden Sea (Germany), Dee Estuary (UK), Kattegat), NW Pacific (Tokyo Bay and Ariake Sound, both in Japan) and the NE Pacific (Vancouver Island, Canada) (Table 2 and Figure 1). We also identify a specimen from the German Bight depicted by Nehring (1997, his figures 23–24, as *Trinovantedinium capitatum*) also as *T. pallidifulum*. The highest relative abundance of *T. pallidifulum* ~~eysts~~ was recorded in a surface sample from the Gulf of Mexico where [it the eysts](#) contributes up to 3.8% of the cyst assemblages (Table 2). *T. pallidifulum* was found in this study in surface sediment samples corresponding to SSTs of ~11.30–31.45 °C and SSSs with a range of 8.60–33.61 psu (Table 2).

**3.5. *Trinovantedinium pallidifulum* cyst wall chemistry**

The spectrum produced by micro-FTIR analysis of the cyst wall (Figure 4) shows: a broad peak centered at 3340 cm<sup>-1</sup> (OH stretching), weak peaks at 2920 and 2860 cm<sup>-1</sup> (aliphatic CH stretching), peaks at 1780 and 1700 cm<sup>-1</sup> (C=O stretching), 1630 cm<sup>-1</sup> (C=C, C=O stretching)

Formatted: Font: Times New Roman, 12 pt

(amide I)), 1585 and 1560  $\text{cm}^{-1}$  (CN stretching, NH bending (amide II)), 1440 and 1405  $\text{cm}^{-1}$  (CH bending), 1350  $\text{cm}^{-1}$  ( $\text{NO}_2$ ,  $\text{CCH}_3$  (not indicated in Figure 4)), 1310  $\text{cm}^{-1}$  (amide III), 1255  $\text{cm}^{-1}$  (NH bending), 1140 and 1030  $\text{cm}^{-1}$  (C-O stretching), and 890, 850 and 770  $\text{cm}^{-1}$  (CH out of plane).

#### 4. Discussion

##### 4.1. Comparative morphology of the motile stage of *P. louisianensis* within the *Conica* group

The motile stage of *P. louisianensis* can be distinguished from all species from the *Conica* group by the unique combination of excavated antapical horns each bearing a single spine, an epitheca with convex sides being longer than the hypotheca, a slightly left-handed cingulum, a stenodeltaform linteloid 2a and plates with polygonal reticulations. A few species show strong similarities. *P. conicum* also has spines on the antapical horns, but has an epitheca that is as long as the hypotheca, straight sutures from the apex to the cingulum and a more elongated pore (Gu et al. 2015). *P. shanghaiense* has no displacement of the cingulum, an epitheca with straight sides, two relatively closer antapical horns that bear no spines and a 2a that was isodeltaform linteloid (Gu et al. 2015). *P. conicoides* also bears spines on the antapical horns, but is more polar compressed. It also has an epitheca that is as long as the hypotheca, a 1''' which has a "nose" at the onset of the sulcus, and straight sutures from the apex to the cingulum (Hoppenrath et al. 2009, p. 158). The plates of *P. obtusum* and *P. leonis* are ornamented with longitudinal ribs (Hoppenrath et al. 2009, p. 158–159).

##### 4.2. Comparative morphology of the cyst of *P. louisianensis* within the *Conica* group

The cyst of *P. louisianensis*, which corresponds to the cyst-based-defined species taxon *Trinovantedinium pallidifulum*, can be easily differentiated from all other species belonging to the genus *Trinovantedinium* (Table 1). The species differs from *Trinovantedinium applanatum* and *Trinovantedinium henrietii* because these are transparent and have an epicyst the same length as the hypocyst (Matsuoka 1987, Louwey et al. 2008). *Trinovantedinium boreale* is also transparent and has processes with platforms (Bujak 1984; see also Head 1994). *Trinovantedinium glorianum* has much more densely distributed and hollow processes and antapical horns that are sharper (Head et al. 1989). *T. variabile*, *T. harpagonium*, *T. ferrugnomatum* and *T. sterthense* are more polar compressed with rounded antapical horns and bear longer processes with distal platforms or taeniate or aculeate processes (Bujak 1984, de Verteuil & Norris 1992, Head 1993). *Trinovantedinium papula* and *Trinovantedinium? xylochoporum* are much more rounded and have longer processes (de Verteuil & Norris 1992).

##### 4.3. Validity of the ~~genus~~ genera *Trinovantedinium* and *Selenopemphix*

The molecular phylogeny shows that *P. louisianensis* and *P. shanghaiense* are most closely related, which is thus the case for their respective cysts, *Trinovantedinium pallidifulum* and *Trinovantedinium applanatum*. This supports is consistent with the monophyly of the genus *Trinovantedinium*. Whether this will be supported by the molecular characterization of the other extant species, *Trinovantedinium variabile*, should be the subject of further study.

The phylogenetic position of *Selenopemphix undulata* is more problematic because it is positioned in another clade than *Protoperidinium conicum*, the motile stage associated with the cyst-based taxon *Selenopemphix quanta* (note that this cyst-theca relationship needs further study, e.g. Matsuoka & Head 2013). This polyphyly suggests that both species belong to two

Formatted: Font: Not Italic

different genera, and their relatively long genetic distance suggests that they diverged a long time ago. This polyphyly is also supported by significant morphological differences between species of both cyst types, particularly the position of the archeopyle, which is more offset in *Selenopemphix undulata* (Verleye et al. 2009) than *Selenopemphix quanta* (it is considered central or slightly offset, see Head 1993, p. 31-32). It is, therefore, likely that *Selenopemphix quanta* should be transferred back to its initial name, *Multispinula quanta*. However, we suggest that the molecular phylogenetic position of the type species of *Selenopemphix*, *Selenopemphix nephroides*, the cyst considered to belong to *Protoperidinium subinerme* (Rochon et al. 1999) needs to be established before such a transfer. Either way, this result emphasizes the importance of the archeopyle position in cyst taxonomy, whereas the reniform cyst shape of both species, seems to be polyphyletic.

**4.4. Validity of the genus Protoperidinium**  
Since long there has been a mismatch between paleontological and biological names, whereas several biological species that belong to the genus *Protoperidinium* have been associated with cysts belonging to different cyst-defined genera (e.g. *Brigantedinium*, *Quinquecuspsis*, *Selenopemphix*, etc.) (e.g. Fensome et al. 1993). Although there have been attempts to reconcile both nomenclatural systems (e.g. Harland 1982), no consensus has yet been reached as how to resolve this issue and here we also suggest to respect the status quo in order to avoid further confusion.

**4.4.4.5. Evolution of Trinovantedinium and Selenopemphix within the Peridinales**  
Previous studies have tried to elucidate evolutionary patterns based on the morphologic changes of the motile stages (e.g., Taylor 1980) or on the basis of cyst morphology (e.g., Bujak & Davies 1983). Molecular phylogenetics largely support the biological approach which focuses on variations in tabulation, and consider the Monovela group and Diplopsaloideans as ancestral to the *Protoperidinium* sensu stricto group (e.g., Liu et al. 2015a,b; Gu et al. 2015; Mertens et al. 2015). Both of these ancestral groups have not been identified in the earlier fossil record of the Peridinales (Cretaceous – Eocene), and the cyst morphologies observed during this time period (Bujak & Davies 1983) are more similar to the cyst-based genera *Lejeunecysta*, *Quinquecuspsis* and *Trinovantedinium*, which, based on molecular phylogenetics, belong to the *Protoperidinium* sensu stricto group (Figure 3). A reinvestigation of the fossil record is urgently needed to explain this discrepancy between the fossil record and the molecular phylogenetics, and begs the question whether preservation issues or the complex cyst identification of these relatively unknown ancestral species could be responsible, or if multigene phylogenies would also support the LSU based phylogeny (e.g., Orr et al. 2012).

*Trinovantedinium boreale* would presumably be the oldest *Trinovantedinium* species, as it has been observed in the late Paleocene, since all other species have Miocene or later first ~~occurrences~~ appearances (Table 1). Two other closely related species to *Trinovantedinium applanatum* and *Trinovantedinium pallidifolium*, the cyst of *Protoperidinium divaricatum* (cyst-defined-based name *Xandarodinium xanthum*) and the cyst of *Protoperidinium leonis* (cyst-defined-based name *Quinquecuspsis concreta*) (Figure 3) both have younger first ~~occurrences~~ appearances (Miocene ~~–~~ Matsuoka, 1992; and Pleistocene ~~–~~ de Vernal et al., 1992), respectively). Interestingly, the here documented *Trinovantedinium applanatum* type B shows similarities to *Trinovantedinium boreale*, although the latter has longer and fewer processes. It

Formatted: Font: Bold

Formatted: Bullets and Numbering

Formatted: Font: Not Italic

would, therefore, be important to obtain a sequence from this type B and see how it relates to the other sequences.

#### **4.5.4.6. Distribution of *Trinovantedinium pallidifulum***

*T. pallidifulum* was shown in this study to be widely distributed in temperate to tropical waters (Figure 1), in surface sediment samples corresponding to SSTs of ~11.30–31.45 °C and SSSs with a range of 8.60–33.61 psu.

#### **4.6.4.7. Cyst wall geochemistry of *Trinovantedinium pallidifulum***

The *T. pallidifulum* cyst wall chemistry demonstrates a composition both consistent and dissimilar with other known heterotrophic dinoflagellates (Figure 4). The greatest similarity is the evidence for nitrogen-containing functional groups in the cyst wall, which is consistent with all previously analyzed cysts from heterotrophic dinoflagellates (Bogus et al. 2014). This evidence includes several absorptions indicative of amide groups (1630, 1585, 1560, 1310, and 1255 cm<sup>-1</sup>). The presence of nitrogen-containing functional groups was suggested to reflect the heterotrophy of the dinoflagellate producing the cyst (Bogus et al. 2014), with the incorporation of prey waste products (high in nitrogen groups from the breakdown of proteinaceous material) into the cyst wall. Additionally, similar to other species, there is clear evidence for alcohol groups (3340, 1140 and 1030 cm<sup>-1</sup>), although the strongest series of absorptions in *T. pallidifulum* is different. In fact, the most obvious difference in the cyst wall chemistry of *T. pallidifulum* compared to other heterotrophic dinoflagellates is the lack of the dominant series of absorptions characteristic for polysaccharides (Bogus et al. 2014; Versteegh et al. 2012). However, absorptions at 1140 and 1030 cm<sup>-1</sup> are present, indicating that sugars may still be present as part of the cyst wall. The strongest absorptions are at 890 and 850 cm<sup>-1</sup>, which usually result from ring vibrations and is supported somewhat by the absorption at 1630 cm<sup>-1</sup>. There is also evidence that this species has a cyst wall that contains a significant carboxylic acid/ester component (1780, 1700 cm<sup>-1</sup>), which may increase the stability of the cyst wall polymer (Yang et al. 1996). This pattern has been suggested in some species of the fossil *Apectodinium* genus (Bogus et al. 2012). However, the cyst wall is not aliphatic because absorptions indicating aliphatic C-H stretching are weak, which is consistent with many dinoflagellate cysts (de Leeuw et al. 2004; Bogus et al. 2014; Versteegh et al. 2012). Therefore, the spectrum of *T. pallidifulum* is unique to any previously analyzed species because it appears to incorporate amide groups, similar to other heterotrophic species, and contain a higher abundance of ester groups, found only in two other fossil species thus far. These results suggest that dinoflagellate cyst wall chemistry may be more diverse than previously considered and further study is required.

## **5. Conclusions**

- We document the cyst-theca relationship for the cyst-based taxon *Trinovantedinium pallidifulum*, and erect the species *Protoperidinium lousianensis* to describe the motile stage.
- LSU rDNA based phylogenies show the closeness of *P. lousianensis* and *P. shanghaiense*, and also of their respective cysts, *Trinovantedinium pallidifulum* and *Trinovantedinium applanatum*.
- The genus *Trinovantedinium* is monophyletic, whilst the genus *Selenopemphix* is polyphyletic.

Formatted: Font: Bold

Formatted: Bullets and Numbering

Formatted: Font: Bold

Formatted: Bullets and Numbering

Formatted: Font: Not Bold



- *Trinovantedinium pallidifulum* is widely distributed, and can be found at SSTs of ~11.30–31.45 °C and SSSs of 8.60–33.61 psu.
- *Trinovantedinium pallidifulum*’s cyst wall chemistry is unique but contains amide groups, which is consistent with other heterotrophic species.

Acknowledgements

Kenneth Neil Mertens is a postdoctoral researcher of FWO Belgium. Martin J. Head is acknowledged for advice on the genus *Trinovantedinium*. Anna Godhe is thanked for providing a sample from the Kattegat. Evelyne Goubert is thanked for providing a sample from La Vilaine Bay and Jean-Pierre Debenay for samples from the Vie River estuary. We acknowledge the Scripps Institution of Oceanography (SIO) Geological Collections and in particular Alex Hangsterfer for sediment samples from the Bay of Bengal. Pieter R. Gurdebeke is thanked for providing a nice specimen of *Trinovantedinium variabile*. K.N.M. and A.M.P. thank Anne de Vernal for letting us use the equipment to conduct germination experiments at GEOTOP. Sofia Ribeiro is thanked for loan of slides from Portugal. G.J.M.V. acknowledges support by the German Science Foundation (Heisenberg fellowship VE486/2). A.M.P. acknowledges the support of the Natural Science and Engineering Research Council of Canada (NSERC). Support for ship time for Gulf of Mexico specimens was provided by the National Oceanic and Atmospheric Administration, Center for Sponsored Coastal Ocean Research, under awards NA06OP0528 and NA09NOS4780204 to Louisiana Universities Marine Consortium and awards NA06OP0529 and NA09NOS4780230 to Louisiana State University. The crew of the R/V *Pelican* is thanked for assistance with sample collection. Martin J. Head and an anonymous reviewer are thanked for remarks that significantly improved the manuscript.

Author biographies

Kenneth Neil Mertens is a ~~postdoctoral permanent~~ researcher at ~~Ghent University, Belgium~~Ifremer, LER BO, Concarneau, France. He received his Ph.D. in 2009 from Ghent University. His research interests are the taxonomy, evolution, phylogeny and biogeography of dinoflagellates, and the palaeoceanographical application of dinoflagellate cysts, particularly in the Quaternary and Neogene.

Haifeng Gu is a Professor at Third Institute of Oceanography, China. He received his Ph.D. in 2007 from Ocean University of China. His research interests are taxonomy and evolution of dinoflagellates.

Yoshihito Takano is a postdoctoral researcher at National Research Institute of Fisheries Science, Fisheries Research Agency, Yokohama, Japan. He received his Ph.D. in 2006 from Hokkaido University. His research interests are taxonomy and evolution of dinoflagellates.

Andrea Price is a PhD candidate in the Department of Geography at McGill University. She recieved her BSc Honours and MSc in Earth and Ocean Sciences at the University of Victoria. Her main research interests include using dinoflagellate cysts as indicators of water quality in North American estuaries, the seasonality of cyst production, and the use of cysts in Quaternary paleoceanographic studies.

Dr. Vera Pospelova is an Associate Professor at the School of Earth and Ocean Sciences (SEOS), University of Victoria. She received her Honours BSc in Geology from the Novosibirsk University (Russia) and PhD in Physical Geography from McGill University (Canada). Vera's research interests include: taxonomy of late Quaternary dinoflagellate cysts; cyst production, ecology, and seasonal dynamics in coastal waters; applications of dinoflagellate cysts as indicators of water quality conditions, with the emphasis on pollution and eutrophication in North American estuaries, as well as paleoceanographic reconstructions using sedimentary archives.

Kara Bogus is a staff scientist with the International Ocean Discovery Program (IODP) at Texas A&M University (College Station, TX, USA). She earned her PhD (marine geosciences) from the University of Bremen (Bremen, Germany). Her research interests include using organic-walled dinoflagellate cysts in paleoceanographic studies, cyst wall chemical composition and its application both to environmental reconstructions and diagenesis of sedimentary organic carbon.

Gerard Versteegh combines organic geochemistry, palynology and palaeobotany to assess organic matter degradation, and, through this, to elucidate present and past environment and evolution. In 1995 received his PhD in Biology at Utrecht University on palynology and the onset of Northern Hemisphere ~~Glaciations~~glaciations. He expanded his expertise with organic geochemistry at the Netherlands Institute for Sea Research. At the Université de Lille 1, France he was an invited Professor in 2007 and 2008 and obtained a prestigious Heisenberg Fellowship from the German Science Foundation in 2009. Currently he is at the Alfred-Wegener-Institute in Bremerhaven and the Centre for Marine Environmental Research, Bremen University. His interests include the macromolecular composition of palynomorphs, selective degradation of organic matter, the Paleozoic terrestrialisation, (sub)recent Mediterranean environmental change, proxy development, lipids, acritarchs and dinoflagellates.

Fabienne Marret is a micropalaeontologist by training, with research interest in the field of Quaternary palaeoceanography and palaeoecology. She received her Ph.D. in 1994 from the University of Bordeaux, France. She studies marine sediments from around the world to interpret past environmental changes, based on vegetation tracers (pollen grains) and sea-surface conditions (dinoflagellate cysts). She is particularly interested in the causes of abrupt climate changes in the past, from the tropics to the poles.

R. Eugene Turner is a faculty member at Louisiana State University. He received his Ph.D. in 1974 from the University of Georgia, USA. His research interests include coastal oceanography and wetlands.

Nancy Rabalais is Executive Director and Professor of the Louisiana Universities Marine Consortium in Chauvin, Louisiana. Since the mid-1980s, Rabalais has been the driving force behind identifying and characterizing the dynamics of the large hypoxic region in the Gulf of Mexico, which receives excess nutrients from the Mississippi River. She serves on numerous boards and panels for federal agencies and national organizations. She has received numerous awards, most recently the John D. & Catherine T. MacArthur 'genius' award (2012). She graduated with a Ph.D. from the University of Texas in 1983.

1  
2  
3  
4  
5  
6  
7  
8  
9 638 Kazumi Matsuoka is a Professor Emeritus at Nagasaki University. ~~After retiring in~~Although  
10 639 retired since 2013, he continues scientific ~~works in~~research on fossil and modern dinoflagellates  
11 640 in particular cyst-motile form relationships of both naked and thecate dinoflagellates.  
12 641

13 642 **References**

14 643  
15 644 Adl SM, Simpson AGB, Farmer MA, Andersen RA, Anderson OR, Barta JR, Bowser SS,  
16 645 Brugerolle G, Fensome RA, Fredericq S, et al. 2005. The new higher level classification of  
17 646 Eukaryotes with emphasis on the taxonomy of Protists. J Eukaryot Microbiol. 52: 399–451.

18 647  
19 648 Artzner DG, Dörhöfer G. 1978. Taxonomic note: *Lejeunecysta* nom. nov. pro *Lejeunia* Gerlach  
20 649 1961 emend. Lentin and Williams 1976 - dinoflagellate cyst genus. Can J Bot 56: 1381–1382.

Formatted: Font: Italic

Formatted: Font: Italic

21 650  
22 651 Baldwin RP. 1987. Dinoflagellate resting cysts isolated from sediments in Marlborough Sounds,  
23 652 New Zealand. New Zeal Mar Fresh. 21: 543–553.

24 653  
25 654 Balech E. 1974. El Genero "*Protoperidinium*" Bergh, 1881 (*Peridinium* Ehrenberg, 1831,  
26 655 partim). Revista Mus Argent Ci Nat, Bernardino Rivadavia Inst Nac Invest Ci Nat Bot. 4: 1–79.

27 656  
28 657 Balech E. 1988. Los dinoflagellados del Atlántico Sudoccidental. Pub Esp Inst Español  
29 658 Oceanografía 1: 1–310.

30 659  
31 660 Benedek PN. 1972. Phytoplanktonen aus dem Mittel- und Oberoligozän von Tönisberg  
32 661 (Niederrheingebiet). Palaeontogr Abt B 137: 1–71.

33 662  
34 663 Bergh RS. 1881. Der Organismus der Cilioflagellaten. Eine phylogenetische Studie. Gegenbaurs  
35 664 Morpholog Jahrb. 7: 177–288.

36 665  
37 666 Boc A, Diallo AB, Makarek V. 2012. T-REX: a web server for inferring, validating and  
38 667 visualizing phylogenetic trees and networks. Nucleic Acids Res. 40: W573–W579.

39 668  
40 669 Boessenkool K, Van Gelder M-J, Brinkhuis H, Troelstra SR. 2001. Distribution of organic-  
41 670 walled dinoflagellate cysts in surface sediments from transects across the Polar Front offshore  
42 671 southeast Greenland. J Quaternary Sci. 16: 661–666.

43 672  
44 673 Bogus K, Harding IC, King A, Charles AK, Zonneveld K, Versteegh, GJM. 2012. The  
45 674 composition of species of the *Apectodinium* complex (Dinoflagellata). Rev Palaeobot Palynol.  
46 675 183: 21–31.

47 676  
48 677 Bogus K, Mertens KNM, Lauwaert J, Harding IC, Vrielinck H, Zonneveld K, Versteegh, GJM.  
49 678 2014. Differences in the chemical composition of organic-walled dinoflagellate resting cysts  
50 679 from phototrophic and heterotrophic dinoflagellates. J Phycol. 50: 254–266.

51 680  
52 681 Bolch CJS. 1997. The use of polytungstate for the separation and concentration of living  
53 682 dinoflagellate cysts from marine sediments. Phycologia 37: 472–478.  
54 683

- Bradford MR. 1977. New species attributable to the dinoflagellate cyst genus *Lejeunia* Gerlach, 1961 emend. Lentin and Williams 1975. Grana 16: 45–59.
- Bringué M, Pospelova V, Pak D. 2013. Seasonal production of organic-walled dinoflagellate cysts in an upwelling system: a sediment trap study from the Santa Barbara Basin, California. Mar Micropaleontol. 100: 34–51.
- Bujak JP. 1984. Cenozoic dinoflagellate cysts and acritarchs from the Bering Sea and northern North Pacific, D.S.D.P. Leg 19. Micropaleontology 30: 180–212.
- Bujak JP, Davies EH. 1983. Modern and fossil Peridiniinae. AASP Contribution Series 13: 1–204.
- Bütschli O. 1885. Erster Band. Protozoa. 3. Unterabtheilung (Ordnung) Dinoflagellata. In: Dr. H.G. Bronn's Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild. C.F. Leipzig und Heidelberg: Winter'sche Verlagshandlung; p. 906–1029.
- Cárdenas G, Cabrera G, Taboada E, Miranda SP. 2004. Chitin characterization by SEM, FTIR, XRD, and <sup>13</sup>C cross polarization/mass angle spinning NMR. J Appl Polym Sci. 93: 1876–1885.
- Cho H-J, Kim C-H, Moon C-H, Matsuoka K. 2003. Dinoflagellate cysts in recent sediments from the southern coastal waters of Korea. Bot Mar. 46: 332–337.
- Colthup NB, Daly LH, Wiberly SE. 1990. Introduction to Infrared and Raman Spectroscopy. London: Academic Press Limited.
- Daugbjerg N, Hansen G, Larsen J, Moestrup Ø. 2000. Phylogeny of some of the major genera of dinoflagellates based on ultrastructure and partial LSU rDNA sequence data, including the erection of three new genera of unarmoured dinoflagellates. Phycologia 39: 302–317.
- de Leeuw JW, Versteegh GJM, van Bergen PF. 2006. Biomacromolecules of algae and plants and their fossil analogues. Plant Ecol. 182: 209–233.
- [De Schepper S, Head MJ, Louwye S. 2009. Pliocene dinoflagellate cyst stratigraphy, palaeoecology and sequence stratigraphy of the Tunnel-Canal Dock, Belgium. Geol Mag 146: 92–112.](#)
- de Vernal A, Londeix L, Mudie PJ, Harland R, Morzadec-Kerfourn M-T, Turon J-L, Wrenn JH. 1992. Quaternary organic-walled dinoflagellate cysts of the North Atlantic Ocean and adjacent seas: ecostratigraphy and biostratigraphy. In: Head MJ, Wrenn JH, editors. Neogene and Quaternary dinoflagellate cysts and acritarchs. Dallas, Texas: American Association of Stratigraphic Palynologists Foundation; p. 289–328.
- de Verteuil L, Norris G. 1992. Miocene Protoperidiniacean dinoflagellate cysts from the Maryland and Virginia coastal plain. In: Head MJ, Wrenn JH, editors. Neogene and Quaternary

1  
2  
3  
4  
5  
6  
7  
8  
9 729 dinoflagellate cysts and acritarchs. Dallas, Texas: American Association of Stratigraphic  
10 730 Palynologists Foundation; p. 391–430.  
11 731  
12 732 Faust MA. 2006. Creation of the subgenus *Testeria* Faust *subgen. nov. Protoperidinium* Bergh  
13 733 from the SW Atlantic Ocean: *Protoperidinium novella* sp. nov. and *Protoperidinium concinna*  
14 734 sp. nov. Dinophyceae. Phycologia 45: 1–9.  
15 735  
16 736 Fensome RA, Taylor FJR, Norris G, Sarjeant WAS, Wharton DI, Williams GL. 1993. A  
17 737 classification of fossil and living dinoflagellates. Micropaleontology Special Publication 7: 1–  
18 738 245.  
19 739  
20 740 Fukuyo Y, Takano H, Chihara M, Matsuoka K. 1990. Red tide organisms in Japan-An illustrated  
21 741 taxonomic guide. In: Fukuyo Y, Takano H, Chihara M, Matsuoka, editors. Tokyo, Japan: Uchida  
22 742 Rokakuho Co. Ltd. (In Japanese/English); 407 p.  
23 743  
24 744 Furio EE, Matsuoka K, Mizushima K, Baula I, Chan KW, Puyong A, Srivilai D, Sidharta, BR,  
25 745 Fukuyo Y. 2006. Assemblage and geographical distribution of dinoflagellate cysts in surface  
26 746 sediments of coastal waters of Sabah, Malaysia. Coastal Marine Science 30: 62–73.  
27 747  
28 748 Gómez F. 2012. A checklist and classification of living dinoflagellates (Dinoflagellata,  
29 749 Alveolata). CICIMAR Océánides 27: 65–140.  
30 750  
31 751 Gribble KE, Anderson DA. 2006. Molecular phylogeny of the heterotrophic dinoflagellates,  
32 752 *Protoperidinium*, *Diplopsalis* and *Preperidinium* (Dinophyceae), inferred from large subunit  
33 753 rDNA. J Phycol 42: 1081–1095.  
34 754  
35 755 Gu H, Liu T, Mertens KN. 2015. Cyst-theca relationship and phylogenetic positions of  
36 756 *Protoperidinium* (Peridinales, Dinophyceae) species of the sections *Conica* and *Tabulata*.  
37 757 Phycologia 54: 49–66.  
38 758  
39 759 Haeckel E. 1894. Systematische Phylogenie. Entwurf eines natürlichen Systems der Organismen  
40 760 auf Grund ihrer Stammesgeschichte. 1. Theil: Systematische Phylogenie der Protisten und  
41 761 Pflanzen: p. I–XV + 1–400, Georg Reimer (Berlin).  
42 762  
43 763 Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis  
44 764 program for Windows 95/98/NT. pp. 95–98.  
45 765  
46 766 Harland R. 1977. Recent and Late Quaternary (Flandrian and Devensian) dinoflagellate cysts  
47 767 from marine continental shelf sediments around the British Isles. Palaeontogr Abt B 164: 87–  
48 768 126.  
49 769  
50 770 Harland R. 1982. A review of recent and quaternary organic-walled dinoflagellate cysts of the  
51 771 genus *Protoperidinium*. Palaeontology 25: 369–397.  
52 772  
53 773  
54 774



- Head MJ, Norris G, Mudie PJ. 1989. 26. New species of dinocysts and a new species of acritarch from the upper Miocene and lowermost Pliocene, ODP Leg 105, Site 646, Labrador Sea. In: Srivastava SP, Arthur M, Clement B, et al., editors. Ocean Drilling Program, Proceedings, Scientific Results, Leg 105, p. 453–466.
- Head MJ. 1993. Dinoflagellates, sporomorphs and other palynomorphs from the Upper Pliocene St. Erth Beds of Cornwall, southwestern England. J Paleo. Memoir 31: 1–62.
- Head MJ. (Ed.) 1994. A forum on Neogene and Quaternary dinoflagellate cysts: the edited transcript of a round table discussion held at the Third Workshop on Neogene and Quaternary Dinoflagellates; with taxonomic appendix. Palynology 17: 201–239. [Imprinted 1993]
- Head MJ, Norris G, Mudie PJ. 1989. 26. New species of dinocysts and a new species of acritarch from the upper Miocene and lowermost Pliocene, ODP Leg 105, Site 646, Labrador Sea. In: Srivastava SP, Arthur M, Clement B, et al., editors. Ocean Drilling Program, Proceedings, Scientific Results, Leg 105, p. 453–466.
- Head MJ, Riding JB, Eidvin T, Chadwick RA. 2004. Palynological and foraminiferal biostratigraphy of (Upper Pliocene) Nordland Group mudstones at Sleipner, northern North Sea. Mar Petrol Geol 21: 277–297.
- Hoppenrath M, Elbrächter M, Drebes G. 2009. Marine Phytoplankton. Selected Microphytoplankton Species from the North Sea around Helgoland and Sylt. Kleine Senckenberg-Reihe 49, 264 pp.
- Jiménez-Moreno G, Head MJ, Harzhauser M. 2006. Early and Middle Miocene dinoflagellate cyst stratigraphy of the Central Paratethys, Central Europe. J Micropal 25: 113–139.
- Jørgensen E. 1912. Bericht über die von der schwedischen Hydrographisch- Biologischen Kommission in den schwedischen Gewässern in den Jahren 1909-1910 eingesammelten Planktonproben. Svenska Hydrogr-Biol Komm Skr. 4: 1–20.
- Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. Nucleic Acids Res. 33: 511–518.
- Kawami H, Matsuoka K. 2009. A new cyst-theca relationship for *Protoperidinium parthenopes* (Peridinales, Dinophyceae). Palynology 33: 11–18.
- Kawami H, van Wezel R, Koeman RPT, Matsuoka K. 2009. *Protoperidinium tricingulatum* sp. nov. (Dinophyceae), a new motile form of a round, brown, and spiny dinoflagellate cyst. Phycol Res. 57: 259–267.
- Kojima N, Seto K, Takayasu K, Nakamura M. 1994. Dinoflagellate cysts assemblage found in the surface sediments of Lake Nakaumi, Western Japan. Laguna 1: 45–51.

1  
2  
3  
4  
5  
6  
7  
8  
9 820 Krepakevich A, Pospelova V. 2010. Tracing the influence of sewage discharge on coastal bays  
10 821 of Southern Vancouver Island (BC, Canada) using sedimentary records of phytoplankton. Cont  
11 822 Shelf Res. 30: 1924–1940.  
12 823  
13 824 Lewis J, Dodge JD, Tett P. 1984. Cyst-theca relationships in some *Protoperidinium* species  
14 825 (Peridinales) from Scottish sea lochs. Journal of Micropalaeontology 3: 25–34.  
15 826  
16 827 Lewis J, Dodge JD. 1987. The cyst-theca relationship of *Protoperidinium americanum* (Gran &  
17 828 Braarud) Balech. Journal of Micropalaeontology 6: 113–121.  
18 829  
19 830 Li Z, Matsuoka K, Shin HH, Kobayashi S, Shin K, Lee T, Han M-S, 2015. *Brigantedinium*  
20 831 *majusculum* is the cyst of *Protoperidinium sinuosum* (Protoperidiniaceae, Dinophyceae).  
21 832 Phycologia 54: 517–529.  
22 833  
23 834 Liu T, Gu H, Mertens KN, Lan D. 2014. A new dinoflagellate species *Protoperidinium*  
24 835 *haizhouense* sp. nov. (Peridinales, Dinophyceae), its cyst-theca relationship and phylogenetic  
25 836 position within the *Monovela* group. Phycol Res. 62: 109–124.  
26 837  
27 838 Liu T, Mertens KN, Ribeiro S, Ellegaard M, Matsuoka H, Gu H. 2015a. Cyst-theca relationships  
28 839 and phylogenetic positions of Peridinales (Dinophyceae) with two anterior intercalary plates,  
29 840 with description of *Archaeoperidinium bailongense* n. sp. and *Protoperidinium fuzhouense* n. sp.  
30 841 Phycol Res. 63: 134–151.  
31 842  
32 843 Liu T, Mertens KN, Gu H. 2015b. Cyst-theca relationship and phylogenetic positions of the  
33 844 diplopsalioideans (Peridinales, Dinophyceae), with description of *Niea* and *Qia* gen. nov.  
34 845 Phycologia 54: 210–232.  
35 846  
36 847 Louwye S, Mertens KN, Vercauteren D. 2008. New dinoflagellate cysts from the Miocene of the  
37 848 Porcupine Basin, offshore southwest Ireland. Palynology 32: 131–142.  
38 849  
39 850 Matsuoka K. 1982. *Protoperidinium pentagonum* (Gran) Balech. In: the working party in  
40 851 taxonomy in the Akashiwo Kenyukai, editors. Synopsis of Red-tide organisms. Sheet no. 108 [In  
41 852 Japanese].  
42 853  
43 854 Matsuoka K. 1987. Organic-walled dinoflagellate cysts from surface sediments of Akkeshi Bay  
44 855 and Lake Saroma, North Japan. Bulletin Faculty of Liberal Arts, Nagasaki University (Natural  
45 856 Science) 28: 35–123.  
46 857  
47 858 Matsuoka K. 1992. Peridiniacean cyst genus *Xandarodinium* in the Miocene Kaminoyama  
48 859 Formation in the western part of Zao Volcano, Yamagata, north Japan. In: Ishizaki K, Saito T.,  
49 860 editors. Centenary of Japanese Micropaleontology. Tokyo, Japan: Terra Scientific Publishing  
50 861 Company; p.449-455.  
51 862  
52 863 Matsuoka K, Head MJ. 2013. Clarifying cyst–motile stage relationships in dinoflagellates. In:  
53 864 Lewis JM, Marret F, Bradley L., editors. Biological and Geological Perspectives of  
54  
55  
56  
57  
58  
59  
60

- Dinoflagellates. London: The Micropalaeontological Society, Special Publications. Geological Society, p. 325–350.
- Matsuoka K, Kawami H, Nagai S, Iwataki M, Takayama H. 2009. Re-examination of cyst-motile relationships of *Polykrikos kofoidii* Chatton and *Polykrikos schwartzii* Bütschli (Gymnodiniales, Dinophyceae). Rev Palaeobot Palynol 154: 79–90.
- Matsuoka K, Joyce LB, Kotani Y, Matsuyama Y. 2003. Modern dinoflagellate cysts in hypertrophic coastal waters of Tokyo Bay, Japan. J Plankton Res. 25: 1461–1470.
- Mertens KN, Yamaguchi A, Kawami H, Ribeiro S, Leander BS, Price AM, Pospelova V, Ellegaard M, Matsuoka K. 2012a. *Archaeoperidinium saanichi* sp. nov.: A new species based on morphological variation of cyst and theca within the *Archaeoperidinium minutum* Jørgensen 1912 species complex. Mar Micropaleontol. 96: 48–62.
- Mertens KN, Bringué M, Van Nieuwenhove N, Takano Y, Pospelova V, Rochon A, de Vernal A, Radi T, Dale B, Patterson RT, Weckström K, Andrén E, Louwye S, Matsuoka K. 2012b. Process length variation of the cyst of the dinoflagellate *Protoceratium reticulatum* in the North Pacific and Baltic-Skagerrak region: calibration as an annual density proxy and first evidence of pseudo-cryptic speciation. J Quaternary Sci. 27: 734–744.
- Mertens KN, Yamaguchi A, Takano Y, Pospelova V, Head MJ, Radi T, Pieńkowski A, de Vernal A, Kawami H, Matsuoka K. 2013. A new heterotrophic dinoflagellate from the Northeastern Pacific, *Protoperidinium fukuyoi*: cyst-theca relationship, phylogeny, distribution and ecology. J Eukaryot Microbiol. 60: 545–563.
- Mertens KN, Takano Y, Yamaguchi A, Gu H, Bogus K, Kremp A, Bagheri S, Matishov G, Matsuoka K. 2015. The molecular characterization of the enigmatic dinoflagellate *Kolkwitzella acuta* reveals an affinity to the *Excentrica* section of the genus *Protoperidinium*. Syst Biodivers. doi:10.1080/14772000.2015.1078855.
- Nehring S. 1997. Dinoflagellate resting cysts from German coastal sediments. Bot Mar. 40: 307–324.
- Orr RJS, Murray SA, Stüken A, Rhodes L, Jakobsen KS. 2012. When naked became armored: An eight-gene phylogeny reveals monophyletic origin of theca in dinoflagellates. PloS one 7: e50004.
- Pascher A. 1914. Über Flagellaten und Algen. Berichte der Deutschen Botanischen Gesellschaft 32: 136–160.
- Posada D. 2008. jModelTest: phylogenetic model averaging. Mol Biol Evol. 25: 1253–1256.
- Pospelova V, Chmura GL, Walker HA. 2004. Environmental factors influencing spatial distribution of dinoflagellate cyst assemblages in shallow lagoons of southern New England (USA). Rev Palaeobot Palynol. 128:7–34.

Formatted: Font: Italic

Formatted: Font: Italic

1  
2  
3  
4  
5  
6  
7  
8  
9 911  
10 912 Pospelova V, de Vernal A, Pedersen TF. 2008. Distribution of dinoflagellate cysts in surface  
11 913 sediments from the northeastern Pacific Ocean (43–25 degrees N) in relation to sea-surface  
12 914 temperature, salinity, productivity and coastal upwelling. Mar Micropaleontol. 68: 21–48.  
13 915  
14 916 Price AM, Pospelova V. 2011. High-resolution sediment trap study of organic-walled  
15 917 dinoflagellate cyst production and biogenic silica flux in Saanich Inlet (BC, Canada). Mar  
16 918 Micropaleontol. 80: 18–43.  
17 919  
18 920 Radi T, de Vernal A. 2004. Dinocyst distribution in surface sediments from the northeastern  
19 921 Pacific margin (40–60°N) in relation to hydrographic conditions, productivity and upwelling.  
20 922 Rev Palaeobot Palynol. 128: 169–193.  
21 923  
22 924 Reid P. 1977. Peridiniacean and glenodiniacean dinoflagellate cysts from the British Isles. Nova  
23 925 Hedwigia 29: 429–463.  
24 926  
25 927 Ribeiro S, Lundholm N, Amorim A, Ellegaard M. 2010. *Protoperidinium minutum*  
26 928 (Dinophyceae) from Portugal: cyst-theca relationship and phylogenetic position on the basis of  
27 929 single-cell SSU and LSU rDNA sequencing. Phycologia 49: 48–63.  
28 930  
29 931 Rochon A, de Vernal A, Turon J-L, Matthiessen J, Head MJ. 1999. Distribution of recent  
30 932 dinoflagellate cysts in surface sediments from the North Atlantic Ocean and adjacent seas in  
31 933 relation to sea-surface parameters. AASP Contribution Series, 35, 146 pp.  
32 934  
33 935 Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed  
34 936 models. Bioinformatics 19: 1572–1574.  
35 937  
36 938 Sarai C, Kawami H, Matsuoka K. 2013. Two new species formally attributed to *Protoperidinium*  
37 939 *oblongum* (Aurivillius) Park et Dodge (Peridinales, Dinophyceae): Evidence from cyst  
38 940 incubation experiments. Rev Palaeobot Palynol. 192: 103–118.  
39 941  
40 942 Scholin CA, Herzog M, Sogin M, Anderson DM. 1994. Identification of group- and strain-  
41 943 specific genetic markers for globally distributed *Alexandrium* (Dinophyceae). II. Sequence  
42 944 analysis of a fragment of the LSU rRNA gene. J Phycol. 30: 999–1011.  
43 945  
44 946 Shin HH, Yoon YH, Kim Y-O, Matsuoka K. 2011. Dinoflagellate Cysts in Surface Sediments  
45 947 from Southern Coast of Korea. Estuaries and Coasts 34: 712–725.  
46 948  
47 949 Soliman A, Ćorić S, Head MJ, Piller W, El Beialy SY. 2012. Lower and Middle Miocene  
48 950 biostratigraphy, Gulf of Suez, Egypt, based on dinoflagellate cysts and calcareous nannofossils.  
49 951 Palynology 36: 1–42.  
50 952  
51 953 Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with  
52 954 thousands of taxa and mixed models. Bioinformatics 22: 2688–2690.  
53 955  
54 956 Taylor F. 1980. On dinoflagellate evolution. BioSystems 13: 65–108.  
55  
56  
57  
58  
59  
60

Formatted: Font: 12 pt


Formatted: Font: 12 pt

Formatted: Font: 12 pt

- 957  
958 Verleye T, Pospelova V, Mertens KN, Louwye S. 2011. The geographical distribution and  
959 (palaeo)ecology of *Selenopemphix undulata* sp. nov., a new late Quaternary dinoflagellate cyst  
960 from the Pacific Ocean. *Mar Micropaleontol.* 78: 65–83.  
961  
962 Versteegh GJM, Blokker P, Bogus K, Harding IC, Lewis J, Oltmanns S, Rochon A, Zonneveld  
963 KAF. 2012. Flash pyrolysis and infrared spectroscopy of cultured and sediment derived  
964 *Lingulodinium polyedrum* (Dinoflagellata) cyst walls. *Org Geochem.* 43: 92–102.  
965  
966 Wall D, Dale B. 1968. Modern dinoflagellate cysts and evolution of the Peridinales.  
967 *Micropaleontology* 14: 265–304.  
968  
969 Wang Z, Matsuoka K, Qi Y, Chen J. 2004. Dinoflagellate cysts in recent sediments from Chinese  
970 coastal waters. *Marine Ecology* 25: 289–311.  
971  
972 Yamaguchi A, Kawamura H, Horiguchi T. 2006. A further phylogenetic study of the  
973 heterotrophic dinoflagellate genus, *Protoperidinium* (Dinophyceae) based on small and large  
974 subunit ribosomal RNA gene sequences. *Phycol Res.* 54: 317–329.  
975  
976 Yamaguchi A, Kawamura H, Horiguchi T. 2007. The phylogenetic position of an unusual  
977 *Protoperidinium* species, *P. bipes* (Peridinales, Dinophyceae), based on small and large subunit  
978 ribosomal RNA gene sequences. *Phycologia* 46: 270–276.  
979  
980 Yamaguchi A, Hoppenrath M, Pospelova V, Horiguchi T, Leander BS. 2011. Molecular  
981 phylogeny of the marine sand-dwelling dinoflagellate *Herdmania litoralis* and an emended  
982 description of the closely related planktonic genus *Archaeoperidinium* Jörgensen. *Eur J Phycol.*  
983 46: 98–112.  
984  
985 Yang CQ, Xu Y, Wang D. 1996. FT-IR spectroscopy study of the polycarboxylic acids used for  
986 paper wet strength improvement. *Ind Eng Chem Res.* 35: 4037–4042.  
987  
988



1  
2  
3  
4  
5  
6  
7  
8  
9 989  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



990 Tables

991  
992 Table 1. List of species belonging to the genus *Trinovantedinium*, their biostratigraphical ranges and dimensions.  
993

Species name	Biostratigraphical range	Length (excluding processes) (µm)	Width (excluding processes) (µm)	Length of processes (µm)	Measurements from
<i>Trinovantedinium applanatum</i> (Bradford 1977) Bujak and Davies 1983	middle Pleistocene (Mudie 1989) - recent (see Rochon et al. 1999 for remarks)	53–87 54–80	47–74 54–80	max 5 max 5–7	Bradford 1977 Reid 1977
<i>Trinovantedinium sterthense</i> Head 1993a	<del>late Pliocene</del> <b>early Pleistocene</b> (Head 1993a)	41(50.0)61	45(49.6)53	4.5(5.5)7.0	Head 1993a
<i>Trinovantedinium variabile</i> (Bujak 1984) de Verteuil and Norris 1992	late Miocene (Bujak 1984) — recent (e.g., Radi & de Vernal 2004; Price & Pospelova 2011)	45–53	44–50	4–6.5	Bujak 1984
<i>Trinovantedinium harpagonium</i> de Verteuil and Norris 1992	<del>middle Miocene (late Miocene (de Verteuil &amp; Norris 1992) Jimenez-Moreno et al. 2006)</del> — late Pliocene (De Schepper et al. 2009)	<del>52(60)78</del>	<del>48(59)68</del>	<del>8–16</del>	de Verteuil & Norris 1992
<i>Trinovantedinium glorianum</i> (Head et al. 1989) de Verteuil and Norris 1992	early Miocene (Louwye et al. 2007) — early Pleistocene (Head et al. 2004)	58(68)73	52(63)70	2–3	de Verteuil & Norris 1992
<i>Trinovantedinium ferrugnomatum</i> de Verteuil and Norris 1992	late Miocene (de Verteuil & Norris 1992) — early Pliocene (De Schepper et al. 2009)	30(39)55	30(39)48	3–6	de Verteuil & Norris 1992
<i>Trinovantedinium papula</i> de Verteuil and Norris 1992	late Miocene (de Verteuil & Norris 1992)	50(65)75	48(57)70	3–6	de Verteuil & Norris 1992
<i>Trinovantedinium pallidifulum</i> Matsuoka 1987	mid Miocene (de Verteuil & Norris 1992 <sup>1</sup> ) — recent (Matsuoka 1987)	52.2–70.8 43.0(58.5)68.4	56.0–63.4 46.3(56.5)63.9	2.5 1.1(1.8)2.7	Matsuoka 1987 This study
<i>Trinovantedinium henrietii</i> Louwye et al. 2008	mid Miocene (Louwye et al. 2008)	81(89)99	61(75)84	5(7)10	Louwye et al. 2008
<i>Trinovantedinium?</i> <i>xylochoporum</i> de Verteuil and Norris 1992	<del>lower Miocene (Soliman et al. 2012) — middle Miocene (Jimenez-Moreno et al., 2006) mid-Miocene (de Verteuil &amp; Norris 1992)</del>	<del>36(51)66</del>	<del>38(45)62</del>	<del>8–20</del>	de Verteuil & Norris 1992
<i>Trinovantedinium boreale</i> Bujak 1984	late Paleocene — late Oligocene (Kurita & Matsuoka 1993; see Head 1994 for discussion)	42–65	42–63	7–15	Bujak 1984

Formatted: French (France)

Formatted: English (U.K.)

Formatted: English (U.S.)

Formatted: English (U.K.)

Formatted: Font: Not Italic

Formatted: English (U.K.)

<sup>1</sup>Our interpretation of the "undefined protoperidiniacean species" depicted in de Verteuil & Norris, plate 2, figs. 9–12.

Table 2. Sampling stations with details on the collection site, relative abundance (%) and how the sample was used in this study.

Station	Location	Latitude (°N)	Longitude (°E)	Water depth (m)	Sampling data	Salinity (psu)	Temperature (°C)	Type of core	Sampled by*	Relative abundance (%)	Notes
Heslwall, Dee Estuary	UK	53,32	-3,12	0	6/02/14	NA	NA	Hand sampling	FM	NA	Germination experiment
Wadden Sea st. 1	Northern Germany	53,72	7,97	0	29/02/12	NA	NA	Hand sampling	GV, KZ, KNM	NA	Germination experiment
Vinga SW	Kattegat	57,55	11,53	77,5	1994	NA	NA	Boxcore	AG	NA	Palynological study
BV3	La Vilaine Bay, France	47,48	-2,44	6	1/04/10	NA	NA	Boxcore	EG	NA	Palynological study
C6B	Gulf of Mexico	28,87	-90,47	18,4	28/07/08	22,17	30,54	Boxcore	GT, NR	0,8	Palynological study
E2A	Gulf of Mexico	28,74	-91,25	15,2	28/07/08	16,14	29,7	Boxcore	GT, NR	0,6	Palynological study
G3	Gulf of Mexico	28,98	-92,00	20,2	28/07/08	21,51	30,45	Boxcore	GT, NR	0,9	Palynological study
J4	Gulf of Mexico	29,29	-93,08	15	25/07/08	27,99	30,52	Boxcore	GT, NR	0,8	Palynological study
A'2	Gulf of Mexico	29,09	-89,50	11,2	27/07/14	8,60	31,45	Boxcore	GT, NR	0,3	Palynological study
A'3	Gulf of Mexico	29,03	-89,53	13,2	27/07/14	13,60	30,99	Boxcore	GT, NR	1,7	Palynological study
A'4	Gulf of Mexico	28,98	-89,57	33,7	27/07/14	15,15	30,90	Boxcore	GT, NR	0,6	Palynological study
A'5	Gulf of Mexico	28,95	-89,58	63,9	27/07/14	17,73	30,19	Boxcore	GT, NR	1,7	Palynological study
A2	Gulf of Mexico	29,24	-89,75	13,5	28/07/14	25,00	30,00	Boxcore	GT, NR	1,2	Palynological study
A5	Gulf of Mexico	29,07	-89,75	30,5	28/07/14	25,30	29,70	Boxcore	GT, NR	1,2	Palynological study
A7	Gulf of Mexico	28,94	-89,75	53,20	28/07/14	24,70	29,90	Boxcore	GT, NR	2,0	Germination experiment / Palynological study
B4	Gulf of Mexico	29,03	-90,12	18,5	28/07/14	28,09	29,55	Boxcore	GT, NR	3,8	Palynological study
C6C	Gulf of Mexico	28,87	-90,49	19,80	28/07/14	23,40	29,40	Boxcore	GT, NR	1,8	Single-cell PCR / Palynological study
D3	Gulf of Mexico	28,72	-90,83	17,8	29/07/14	15,05	30,83	Boxcore	GT, NR	2,4	Palynological study
F0	Gulf of Mexico	29,27	-91,62	8,0	31/07/14	23,20	31,30	Boxcore	GT, NR	3,7	Palynological study
F3	Gulf of Mexico	28,88	-91,62	19,9	30/07/14	29,15	29,92	Boxcore	GT, NR	1,5	Palynological study
F6	Gulf of Mexico	28,58	-91,62	39,6	30/07/14	30,02	29,51	Boxcore	GT, NR	0,8	Palynological study
I4	Gulf of Mexico	29,18	-92,75	20,8	1/08/14	32,40	30,00	Boxcore	GT, NR	0,3	Palynological study
K4	Gulf of Mexico	29,33	-93,42	17,3	1/08/14	31,74	29,92	Boxcore	GT, NR	0,8	Palynological study
P6	Gulf of Mexico	29,00	-93,71	20,0	1/08/14	33,61	30,08	Boxcore	GT, NR	0,3	Palynological study
SR8	Lake Saroma	44,09	143,52	ca. 7	July 1980	33-33.2	0-15	TFO corer	YF	0,5	Palynological study

AK2	Akkeshi Bay, Japan	43,02	144,47	ca. 6	July 1980	32.6-33	1.5(winter)- 14(summer)	TFO corer	YF	0.5-1.8	Palynological study
AB40	Ariake Sound, Japan	33,08	130,28	10,9	7/06/05	20.0- 30.2	9.2-28.5	TFO corer	KM	NA	Palynological study
Tkb8	Tokyo Bay, Japan	35,27	139,45	ca.12	1/08/99	26.8- 30.5	10.4-30.1	KK type corer	RV	NA	Palynological study
Nagayo-Ura	Omura Bay, Japan	32,51	129,52	11	June 2004	30.8- 33.4	9.5-30.5	sediment trap	HK, KM	NA	Palynological study
YJ1	Yeoja Bay, Korea	34,42	127,31	11	May 2006	NA	NA	Scuba diver	KM, HC	NA	Palynological study
D1	Deukryang Bay, Korea	34,40	127,11	10	1/04/01	NA	NA	Gravity corer	HC	NA	Palynological study
Casino coast	Brazil	-32,12	-52,10	ca.10	7/03/97	NA	NA	TFO corer	KM	NA	Palynological study
Saanich Inlet st. S13	Vancouver Island, BC, Canada	48,75	-123,61	60	24/05/07	28,5	11,3	Petite Ponar Grab	VP	NA	Palynological study

\* AG = Anna Godhe, EG = Evelyne Goubert, FM = Fabienne Marret, GV= Gerard Versteegh, GT = Gene Turner, HC = Hyun-Jin Cho, HK = Hisae Kawami, KM = Kazumi Matsuoka, KNM = Kenneth Neil Mertens, KZ = Karin Zonneveld, NR = Nancy Rabalais, RV = R/V Shirafuji-Maru, VP = Vera Pospelova, YF = Yasuo Fukuyo

1000  
1001

Supplementary table 1. Sampling stations with details on latitude, longitude and water depth.

Name of station	Locality	Latitude (°N)	Longitude (°E)	Water depth (m)	Reference
Akkeshi Bay AK1	Hokkaido, Japan	43,04	144,83	25	Matsuoka 1987
Akkeshi Bay AK2	Hokkaido, Japan	43,02	144,79	25	Matsuoka 1987
Akkeshi Bay AK3	Hokkaido, Japan	42,99	144,79	25	Matsuoka 1987
Akkeshi Bay AK4	Hokkaido, Japan	42,95	144,86	25	Matsuoka 1987
Saroma lake SR8	Hokkaido, Japan	44,14	143,86	16	Matsuoka 1987
6	Lake Nakaumi, Japan	35,52	134,08	NA	Kojima et al. 1994
8	Lake Nakaumi, Japan	35,52	134,08	NA	Kojima et al. 1994
24	Lake Nakaumi, Japan	35,52	134,08	NA	Kojima et al. 1994
25	Lake Nakaumi, Japan	35,52	134,08	NA	Kojima et al. 1994
26	Lake Nakaumi, Japan	35,52	134,08	NA	Kojima et al. 1994
TKB-6	Tokyo Bay, Japan	35,50	139,83	NA	Matsuoka et al. 2003
D1	South of South Korea	34,67	127,18	10	Cho et al. 2003
4	Chinese coast	22,56	114,51	6	Wang et al. 2004
Kuala Penyu Lagoon 1	Sabah, Malaysia	5,53	115,62	NA	Furio et al. 2006
Kuala Penyu Lagoon 2	Sabah, Malaysia	5,57	115,60	NA	Furio et al. 2006
YJB1	Yeoja Bay, South Korea	34,78	127,50	NA	Shin et al. 2011
YJB2	Yeoja Bay, South Korea	34,74	127,50	NA	Shin et al. 2011
YJB3	Yeoja Bay, South Korea	34,71	127,50	NA	Shin et al. 2011
YJB4	Yeoja Bay, South Korea	34,67	127,51	NA	Shin et al. 2011
YJB6	Yeoja Bay, South Korea	34,50	127,59	NA	Shin et al. 2011
YJB8	Yeoja Bay, South Korea	34,43	127,70	NA	Shin et al. 2011
GMB3	Gamak Bay, South Korea	34,60	127,70	NA	Shin et al. 2011



## Figure captions

Figure 1. Map showing locations where *Trinovantedinium pallidifulum* was found in this study (red circles) and in previous studies (blue circles) (see Table 1 and Supplementary Table 1 for respective sample localities).

Figure 2. Drawings of *Protoperidinium louisianensis* n. sp. A). Ventral view. B). Dorsal view.

Figure 3. Molecular phylogeny. A phylogenetic tree using Bayesian inference inferred from LSU rDNA sequences based on new sequence data for *Protoperidinium louisianensis* n. sp. from the Gulf of Mexico and *Selenopemphix undulata* from Brentwood Bay, BC, Canada. The ML bootstrap support values (ML) over 50 and Bayesian posterior probabilities (PP) over 0.7 are shown at the nodes (ML/PP). The black circles indicate maximal support. Clades are labeled and marked with vertical lines on the right, with dashed lines indicating sections of the *Protoperidinium sensu stricto* clade. The scale bar represents inferred evolutionary distance in changes/site. New sequences obtained in this study are indicated in bold font.


Figure 4. *Trinovantedinium pallidifulum* (German Wadden Sea) cyst wall FTIR spectrum in comparison with other cysts produced by heterotrophic dinoflagellates. The spectra from *D. caperatum*, *Brigantedinium* sp., cysts of *Polykrikos- kofoidii sensu Matsuoka et al. (2009)* and *Polykrikos- schwartzii sensu Matsuoka et al. (2009)* are modified from Bogus et al. (2014).

## Plate captions

Plate 1. Figures 1–9. Different views of single specimen of motile stage of *Protoperidinium louisianensis* germinated from cyst depicted in Figures 10–15, and isolated from the Northern Gulf of Mexico. 1. General shape of cell. 2. Ventral view. 3. Lateral view. 4. Dorsal view showing the second intercalary plate (2a). 5. Lateral view. 6. Lateral view. 7. Ventral view showing sulcal area. 8. Dorsal view. 9. Ventral view of hypotheca. Figures 10–15. Cyst stage of *Protoperidinium louisianensis*, corresponding to *Trinovantedinium pallidifulum*, with motile stage depicted in Figures 1–9. 10. Ventral view. 11. Cross section. 12. Dorsal view showing operculum. 13. Peritabular distribution of processes. 14–15. Ventral view showing process distribution on the hypotheca. All scale bars = 10 µm.

Plate 2. Figures 1–9. Motile stage of *Protoperidinium louisianensis* germinated from cyst depicted in Figures 10–15, isolated from the Northern Gulf of Mexico. 1. Cross section showing general shape and cell contents. 2. Ventral view. 3. Ventral view showing shape of the first apical plate. 4. Focus on the apical pore. 5. Dorsal view showing shape of second intercalary plate (2a). 6. Lateral view. 7. Ventral view showing sulcal area. 8. Dorsal view. 9. View of dorsal side of hypotheca. Figures 10–15. Cyst stage of *Protoperidinium louisianensis*, corresponding to *Trinovantedinium pallidifulum*, with motile stage depicted in Figures 1–9. 10. General shape of cyst. 11. Ventral view showing peritabular distribution of processes. 12. Ventral view showing presence of 2 flagellar scars. 13. Lateral view. 14. Dorsal view showing operculum. 15. Dorsal view of hypotheca showing presence of striations on this specimen. All scale bars = 10 µm.

1  
2  
3  
4  
5  
6  
7  
8  
9 1047 Plate 3. Figures 1–12. *Trinovantedinium pallidifulum* from German Wadden Sea. 1–3. Cyst  
10 1048 with cell contents from St. 1. 4–8. Different orientations of germinated specimen. 9–12. Other  
11 1049 specimen). 9. Apical pore. Figures 13–16. Holotype of *Trinovantedinium pallidifulum*. Slide no.  
12 1050 | AK2–2 (87.4/27.6); Sample no. AK2, ~~Recent~~ modern sediment in Akkeshi Bay, Hokkaido,  
13 1051 North Japan (Matsuoka 1987). All scale bars = 10 µm.  
14 1052  
15 1053 Plate 4. Figures 1–12. *Trinovantedinium applanatum*. 1–4. High focus to low focus of single  
16 1054 specimen from 11–12 cm depth in core po287-39-1B (37.75°N, 8.87°W, 92 m water depth, Mira,  
17 1055 Portugal). 5–6, 8–10. Different orientations of single specimen isolated from the German  
18 1056 Wadden Sea (location shown in Table 2). 7. Other specimen from the German Wadden Sea. 11–  
19 1057 13. Scanning electron microscope photographs of specimens isolated from Vie River st. 10  
20 1058 (46.70°N, 1.94°W, France). All scale bars = 10 µm.  
21 1059  
22 1060 Plate 5. *Trinovantedinium applanatum* type A and B occurring in recent sediments. 1–2, 5–7.  
23 1061 Different views of single specimen from Type A from station 1 (32.93°N, 129.86°E, 11.1 m  
24 1062 water depth, Omura Bay, Japan). 3. Type A from Red Sea (va01-200P, 0-5 cm depth, 16.67°N,  
25 1063 41.32°E, , 84 m water depth). 4. Specimen from 1–2 cm core depth from core CIRCE03AR 25P  
26 1064 (15.30°N, 83.39°E, 3145m water depth, Bay of Bengal). 8–9. Type B from sample 11B  
27 1065 (35.87°W, 64.11°N, 1318 m water depth, offshore Greenland, Boessenkool et al. 2001). 10. Type  
28 1066 B from sample 1B (60.02°N, 11.76°W, water depth unknown, offshore Greenland, Boessenkool  
29 1067 et al. 2001). All scale bars = 10 µm.  
30 1068  
31 1069 Plate 6. 1–11. *Trinovantedinium variabile*. 1–6. Specimen from Saanich inlet (UVic 13-303-1,  
32 1070 48.59°N, 123.50°W, water depth 226 m). 7–9. Other specimen from sediment-trap from Saanich  
33 1071 Inlet (UVic 09-183, slide 1, 48.65°N, 123.48°W, water depth 96 m), previously studied by Price  
34 1072 and Pospelova (2011). 10–11. Scanning electron microscope photographs of two specimens from  
35 1073 sample Exp37A from Saanich Inlet (48.55°N, 123.53W, water depth 215 m). Figures 12–14.  
36 1074 *Selenopemphix undulata* from Brentwood Bay, BC, Canada, sequenced through single-cell PCR.  
37 1075 All scale bars = 10 µm.  
38 1076  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60









Figure 2. Drawings of *Protoperidinium lousianensis* n. sp. A). Ventral view. B). Dorsal view.  
55x33mm (300 x 300 DPI)











209x258mm (300 x 300 DPI)




209x267mm (300 x 300 DPI)




209x297mm (300 x 300 DPI)






209x225mm (300 x 300 DPI)



209x191mm (300 x 300 DPI)



209x297mm (300 x 300 DPI)